

Journal of Police Medicine



ORIGINAL ARTICLE

OPEN 6 ACCESS

Descriptive and analytical study of factors affecting life expectancy in doctors on duty in the emergency department

Shabnam Toloei^{1 MSc}, Hamed Aghdam^{2*MD}, Omid Shirzad^{3 PhD}, Pouya Vakilipour^{2 MD}

- ¹ Department of Educational Psychology, Faculty of Humanities, Islamic Azad University, Arak Branch, Arak, Iran.
- ² Research Center for Trauma in Police Operations, Directorate of Health, Rescue & Treatment, Police Headquarter, Tehran, Iran.
- ³ Department of Law, Faculty of Law, Political Sciences and Foreign Languages, Mashhad Branch, Islamic Azad University, Mashhad, Iran.

ABSTRACT

AIMS: Shift work is implemented in various formats or systems, each with its characteristics, advantages, and disadvantages, and can adversely affect various aspects of human life. Reduced life expectancy affects physical and mental health and is the final consequence of job stress in many shift work occupations, including nursing. This study aims to investigate the factors affecting life expectancy in shift doctors in the emergency department.

MATERIALS AND METHODS: This is a descriptive-analytical and cross-sectional study conducted in 2023 at the Applied Research Center of Police in which shift-working doctors of the emergency department were included. The study population consisted of shift doctors in the emergency department of 50 hospitals and clinics related to Iran and Alborz Universities of Tehran and Karaj with at least 6 months of continuous work experience with fixed and rotating shifts in 2022. Sampling was done using a stratified method with a sample size of 300 people (95% confidence level). For this purpose, Miller's Hope Questionnaire along with a questionnaire to collect demographic information was provided to the doctors in different work shifts while explaining the objectives of the study and training instructions and how to complete them. Then the data was analyzed by SPSS software version 26. Appropriate parametric and nonparametric tests, including t-test and ANOVA, were utilized to compare numeric data. Also, correlations between numeric data were assessed through Pearson's correlation and linear regression.

FINDINGS: 313 people (158 women), including 94 general practitioners (30%), 197 emergency physicians (62.9%), and 22 physicians specializing in other fields (7%), with a mean age of 44.07 ± 7.47 years, were included in the study. The mean score of the life expectancy test had significant differences in different subgroups of marital status (p<0.001), parenthood (p<0.001), child living with the parent in question (p<0.001), field of study (p<0.001), spouse employment (p=0.008), level of satisfaction with one's work situation (p<0.001), level of satisfaction with one's life situation (p<0.001), desire to change careers in the current situation (p<0.001), traveling to relieve job stress (p<0.001), interacting with relatives to relieve job stress (p<0.001), exercising to relieve job stress (p<0.001), adequate sleep during the day (p<0.001), specific illness of first-degree family members (p=0.04), and shift status in the emergency room (p<0.001). Also, the results of linear regression showed that the age of physicians, the duration of physicians' marriage, the overall work experience of physicians, and the experience of physicians working in the emergency department significantly predicted the life expectancy test score (R2=0.216 and F=(5,211)11.64 and p<0.001).

CONCLUSION: Inappropriate shift work reduces life expectancy and, as a result, disrupts the personal, family, and professional lives of healthcare personnel. In addition, demographic characteristics, shift work, adequate nighttime sleep, living conditions, travel, exercise, contact with relatives, specific physical or psychological illnesses, and specific illnesses of family members have a significant effect on life expectancy. Also, increasing shift work of emergency department physicians, especially night shifts, significantly reduces the life expectancy of this group of physicians.

KEYWORDS: Shift Work Schedule; Psychometrics; Personal Satisfaction; Hope; Emergency Service, Hospital.

How to cite this article:

Toloei S, Aghdam H, Shirzad O, Vakilipour P. Descriptive and analytical study of factors affecting life expectancy in doctors on duty in the emergency department. I Police Med. 2024;13:e13.

*Correspondence:

Address: Applied Research Center, 5th floor, Vali asr Hospital, Valiasr street, Tehran, Iran, Postal code: 1417944661

Mail: hamedaghdam@gmail.com

Article History:

Received: 05/05/2023 Accepted: 19/07/2024 ePublished: 24/07/2024

Copyright © 2024, Journal of Police Medicine | This open access article is licensed under a Creative Commons Attribution-NonCommercial 4.0 International License

Descriptive and analytical study of factors affecting life expectancy in doctors on duty in the emergency department

INTRODUCTION

Shift work is a social phenomenon that has its roots in human history. Today, for economic and technological reasons, shift work systems have become part of the 24-hour society worldwide. This is because the division of labor must be carried out in such a way that active people are present at the workplace at all hours of the day and night, and it is implemented in many industries and industrial-manufacturing factories, as well as in some service occupations such as medicine, nursing, firefighting, police, water services, electricity, telephone, etc. [1]. There are various formats or systems of shift work, each of which has its characteristics, advantages, and disadvantages. The two general shift work systems are the 12-hour shift system (two shifts) and the 8-hour system (three shifts). A general definition of shift work is an unusual work pattern compared to day work, which includes several types of shift work, including fixed shifts, in which some people work consistently in the morning, afternoon, or night shifts. Rotating shifts, in which people work in rotation in the morning, afternoon, and night shifts, and include forward (clockwise) rotational shifts and backward rotational shifts, which are the opposite of the first and can be considered as fast rotation (2 to 3 days), medium rotation (1 week), and slow rotation (2 to 3 weeks) for each work shift [2]. Because shift workers are forced to be awake and active both during the day when society is active and at night, when society is asleep, this phenomenon is now considered a health threat that can have adverse effects on various aspects of human life [3].

Several studies have been conducted to investigate the negative effects of shift work on the performance, health, and quality of life of shift workers, some of which have shown a relationship between shift work and sleep disorders, gastrointestinal problems, cardiovascular diseases, and mental disorders [4].

Shift work has a long history in the medical profession, in health care institutions and centers, and especially in the emergency department, and has long been a common occurrence. Since acute illnesses, the instability of the physical and mental state, and the dependence of patients on others, require the provision of continuous and ongoing services. Therefore, the nature of such services is far from being planned and people must be present continuously and around the clock to provide services. Based on studies, the prevalence of shift work in European countries is 15 to 20%, and in the United States 20% of the workforce [5]. This prevalence rate would be a good reason to study and investigate the negative effects on the health,

safety, and efficiency of workers. Shift work is one of the oldest challenges for healthcare workers and can have adverse physiological, psychological, and social effects on shift workers [6].

Social, psychological, and family consequences include irritability, depression, stress, and disruption of social relationships, among other psychological phenomena resulting from the interference of shift work with individuals' social and family lives. Shift work can have a negative impact on individuals' family lives and disrupt their social lives, especially when the shift work schedule is such that the individual is away from home in the late afternoon and evening [6].

home in the late afternoon and evening [6]. In developed countries, numerous studies have been conducted on the effects of long working hours in 12-hour shifts and one week in different occupational groups, as well as the consequences of working in 12-hour shifts and 2-week periods, and in which the effects of long working hours and shift work schedules on performance, health, quality of life, and occupational accidents have been explored [7], the results of these studies cannot provide a completely appropriate practical model for our country due to organizational, environmental, and cultural differences. Also, based on studies conducted on shift work schedules, it has been found that a poorly designed work schedule can affect the quality of health care, patient satisfaction, length of stay, absenteeism, cost-effectiveness, and productivity [8]. Most of these studies have been conducted on nursing staff and other people working in healthcare wards, and rarely on physicians. In recent years, with the expansion of the specialty of emergency medicine and the requirement for medical centers to have emergency medicine specialists in the emergency department, this issue has become more necessary to study. In outpatient medical centers and hospital wards, general practitioners are also present, and it is also necessary to study the effect of shift work on this group. Hope and hope for life are concepts that are related to adaptation, faith, and empowerment, and can be considered as an effective response to stressors by making existing conditions bearable. Hope is increased by successful life experiences and decreased by failure experiences. Those with a positive explanatory style have greater happiness, hope, and life satisfaction [9]. Hope is an empowering, multidimensional, dynamic, and powerful factor in adapting to problems and even inconsistencies. Hope is one of the basic concepts of positive psychology and an energizing factor in people's lives. People with higher hopes have more power to achieve their goals, and when they encounter an obstacle, they can maintain their motivation

Toloei et al.

and use alternative ways; however, frustrated people lose their motivation when faced with obstacles because they have few options. Hope is also defined as a psychological energy focused on a goal and a sense of determination to achieve a goal. People with high strategic hopes consider multiple ways to deal with possible obstacles. In contrast, individuals with low strategic hope are less likely to find a feasible solution to achieve their goals. Individuals who can manage conflict in relationships by using positive methods and less negative interactions create an environment in which there is more opportunity for selfdisclosure and agreement on life problems [10]. Schneider et al. have considered hope to be a two-dimensional construct: one is the ability to design paths to desired goals despite obstacles (strategic thinking) and the other is the ability to generate motivation (agentic thinking) to initiate and continue moving along these paths [11]. They described hope as a cognitive complex that is based on a sense of success, arising from various sources (purposeful decisions) and paths (the chosen method to achieve goals). Individuals with high hopes have a sense of goal-setting management, along with the ability to generate plans to achieve their goals. Schneider and Michael also suggested that hope may have a moderating effect. Thus, people with high hopes are more capable of implementing strategies for coping with crises than those with low hopes. Those with more hope can conceptualize their goals, feel more confident and capable of effectively managing stressors, and know that they must have a specific plan to deal with any stress [12]. Life expectancy is one of the fundamental variables that can help healthcare professionals adapt to specific working conditions, adapt to the stressful conditions of the emergency room, and comply with scientific and administrative systems. Hope is one of the most important topics in positive psychology. Positive psychologists believe that hope can protect people from stressful life events. Hope includes a person's perceptions of their abilities to set clear goals, develop plans to achieve these goals, and find and maintain the energy and motivation needed to pursue and follow through on those goals. Hope includes a person's vision of a positive future and increases their efforts to improve their circumstances. There is evidence that high levels of hope are associated with success in various activities, quality of life, and better physical and

Hope plays an important role in effectively dealing with problems and the stress they cause, and is a potentially powerful factor in patients' recovery and adaptation. Therefore, hope is associated

mental health [13].

with increased purpose in personal and work life [14]. In a study by *Cao et al.* [15], they showed that higher hope in medical staff predicts their better participation in the treatment process. Also, *Clayton* et al., in a meta-analysis study, showed that in general, a sense of hope is an important factor for motivation in the treatment process and that people with higher life expectancy have better satisfaction [16].

According to previous studies, it can be seen that shift work, by causing mental and psychological problems in individuals, can weaken their hope, which in the case of doctors, can have harmful effects on patients and the country's health care system. Therefore, this study was conducted to investigate the factors affecting life expectancy in shift doctors working in the emergency department.

MATERIALS & METHODS

This study was descriptive-analytical. The study population was shift physicians working in 50 hospitals and clinic centers affiliated with Iran and Alborz Universities of Medical Sciences, which included a total of 1000 people. The study subjects were selected from among physicians with at least 6 months of continuous professional experience, with fixed and rotating shifts in 2022. According to the studies conducted, the number of physicians working in the emergency departments of the study centers was similar. Therefore, a simple random sampling method was used. For this purpose, first, a list of all physicians working in the emergency departments in all centers was prepared. Then, each physician was assigned a specific code for random selection. Finally, using Excel software, the required number of random codes was determined and the study samples were selected. The sample size required for the study was calculated as 285 people, considering a confidence level of 95% and a z value of 1.96, a corrected permissible sampling error of 5.8%, a predicted success rate (p) of 0.5, and using the formula:

 $E2-1)/p\times(p\times Z2=n$

Considering the probability of statistical attrition of 300 people ,a larger number 350) items (of questionnaires were distributed.

A questionnaire consisting of two sections of demographic information and a life expectancy questionnaire) HOL (was used as a data collection tool. The questions on demographic characteristics and personal characteristics included gender, age, and work history, work history in the emergency department, work shift, income, sleep, living situation, travel, exercise, and interaction with relatives, specific illness, and family members

Descriptive and analytical study of factors affecting life expectancy in doctors on duty in the emergency department

in different occupational groups of emergency department physicians.

To measure the level of hope in the study subjects, the Miller Hope Questionnaire or Miller Hope Questionnaire) MHS (was used . This questionnaire is the most common tool for measuring life expectancy ,which consists of 48 separate items on feelings and attitudes and measures different aspects of life expectancy . This questionnaire is translated from an English-language instrument that has not been implemented in Iran so far and requires validation) validity and reliability measurement .(Miller reported the validity of this questionnaire as desirable and its reliability according to Cronbach's alpha was above 80 [17]. The responses to this questionnaire are arranged on a five-point continuum (strongly disagree = 1, disagree = 2, indifferent = 3, agree = 4, strongly agree = 5). In the Miller test, the range of points obtained varies from 48 to 240. 12 items in the Miller questionnaire consist of negative items, and these numbers are scored inversely in evaluation and scoring [17].

A pilot test of this measure was conducted on 75 students in the United States. This test examined the items of hope in terms of their level of readiness and validity. The reliability coefficient was 95%, the validity (retest with a two-week interval) was 87%, and the average home was 175.6 [18]. This test was first administered in Iran by Samiei (1989) on 30 male nursing students, and to determine its reliability, the retest method was used with a oneweek interval. The Pearson correlation coefficient between the two test scores was 75%, and its average was 151.3. To ensure the reliability of the test, this questionnaire was administered to a larger number (152 people) of male and female students of teacher training centers in Tehran, and the Cronbach's alpha coefficient was 91.5%, and the average was 196 [19].

After obtaining permission, to complete the questionnaires and collect data, the subjects visited the clinic and hospital during different work shifts. The objectives of the study were explained, the instructions were explained, and the questionnaire was provided online to the physicians who had agreed to cooperate. All questions in the questionnaire were completed by the subjects in a self-reported manner. In this study, the life expectancy was first determined, and then the average life expectancy score among the physicians and different work shifts was obtained. The information obtained above was analyzed based on the demographic information of the subjects and questions related to possible causes affecting life expectancy.

Ethical Permissions: This study was conducted

by the ethical principles outlined in the Declaration of Helsinki. Ethical approval was obtained from the Ethics Committee for Biomedical Research of the Director of Health Police with the ethics code IR.SBMU.TEB.POLICE.REC.1402.058. All participants provided written informed consent before entering the study. Participants were assured of their right to withdraw from the study at any time without any consequences. All data collected were anonymized to protect the confidentiality of the participants.

Statistical Analysis: The normality of the distribution of quantitative data was examined using the Kolmogorov-Smirnov test. Parametric tests, including t-test and ANOVA, were used to compare data with normal distribution, and nonparametric tests were used to compare data with non-normal distribution. Pearson correlation and stepwise linear T-regression were used to examine the relationship between quantitative variables. In all statistical analyses, a p-value level of less than 0.05 was considered significant. All statistical analyses were performed using SPSS 26 software.

FINDINGS

According to the inclusion criteria, 313 physicians, including 94 general practitioners (30.0%), 197 emergency physicians (62.9%), and 22 physicians specializing in other fields (7.0%), were included in the study. 158 were female (50.50%) and 155 were male (49.50%) with a mean age of 44.07±7.47 years, of which 69.3% were married (30.7% were single). Also, 50.5% of the participants in the study had children, and among those with children, 76.3% lived with their children. 91.3% of the study subjects had a working spouse.

The results of the health status survey revealed that 42.8% of the study subjects suffered from physical or mental illness. Also, 35.8% of the subjects reported a history of a specific illness in their first-degree relatives. In the study of the type of shift work, most people worked in rotating shifts (79.9%). Fixed night shifts were the most common among the types of fixed shifts (about 66%). In the study of the number of shifts per month, 16.6% had less than 4 shifts per month, 55.9% had between 4 and 8 shifts per month, and 27.5% had more than 8 shifts per month. Also, 97.4% of people had a day off shift in the past year. In the study of the salary-work ratio, 86.6% of people believed that their salary was not proportional to their work. Details of the demographic characteristics and employment status of the people are shown in Table 1.

Also, in the study of the level of life satisfaction, 7.3% of people were highly satisfied, and the level

Toloei et al.

of job satisfaction was 38.3% very low, 33.9% low, and 27.8% average. None of the participants in this study were very satisfied with their jobs. In the study of people's desire to change jobs, 34.2% had a very high desire, 33.5% a high desire, 15.3% a moderate desire, 5.8% a low desire, and 11.2% a very low desire to change jobs. In the study of factors to relieve job stress, a small number of people had a high and very high desire to travel, exercise, and spend time with loved ones to reduce job stress (6.7%, 6.4%, and 14.1%, respectively). In the study of nighttime sleep, about 23.6% of people reported adequate sleep during the night most of the time, and 8.6% of people did not have adequate sleep at all. The details of the questionnaire results are given in the form of a Likert column chart in Figure 1.

Table 1) Demographic characteristics and employment status of the study subjects

Row	Variable	Status	Number	Percentage
1	Sex	Female	158	50.5
1	sex	Male	155	49.5
		General Practitioner	94	30.0
2	Educational field	Emergency Medicine Specialist	197	62.9
		Specialist in Other Fields	22	7.0
3	Marital status	Married	217	69.3
	Mai itai Status	Single	96	30.7
4	Having a shild	Yes	158	50.5
4	Having a child	No	155	49.5

5	Child living with the	Yes	158	76.3
5	parent in question	No	49	23.7
6	Whether the spouse	Yes	209	91.3
	is employed	No	20	8.7
		PhD	170	70.5
7	Spouse's latest educational qualification	Master's Degree	12	5.0
	quanication	Bachelor's Degree	59	24.5
8	Specific physical or	Yes	134	42.8
	psychological illness	No	179	57.2
9	Specific illness of first-degree family	Yes	112	35.8
	members	No	201	64.2
10	Average income ratio to activity	Yes	42	13.4
	to activity	No	271	86.6
	Type of shift in	Fixed Shift	63	20.1
11	the emergency department	Rotating Shift	250	79.9
		Morning	19	6.1
		Evening	1	0.3
	Time of shift in	Night	35	11.2
12	the emergency department	Combined Morning, Evening, and Night Shifts	258	82.4
13	Holiday shift in the	Yes	305	97.4
13	past year	No	8	2.6
		Less than 4 nights	52	16.6
14	Number of night shifts in the past year	to 8 nights 5	175	55.9
	omito in the past year	More than 8 nights	86	27.5




Figure 1) Likert chart of the results of the questionnaire taken from the study subjects

The results of the Kolmogorov-Smirnov test showed that the data of the variables of age, duration of marriage, total work history, work history in the emergency department, and life expectancy questionnaire score have a normal distribution. Therefore, to compare the mean score of the life expectancy questionnaire, the Sample

t-test or ANOVA test was used between different groups. The results showed that the mean score of the life expectancy questionnaire in different subgroups of marital status, parenthood, child living with the parent in question, field of study, spouse employment, child living with the parent in question, field of study, spouse employment,

6 Descriptive and analytical study of factors affecting life expectancy in doctors on duty in the emergency department

satisfaction with one's work situation, satisfaction with one's life situation, desire to change career in the current situation, traveling to relieve job stress, interacting with relatives to relieve job stress, exercising to relieve job stress, adequate sleep during the day, specific illness of first-degree family members, and shift status in the emergency department, had a significant difference.

However, the mean score of the life expectancy questionnaire did not differ significantly across different subgroups of gender, spouse's latest educational qualification, average income-activity ratio, specific physical or psychological illness, shift status in the emergency department, day-off shift in the past year, and number of night shifts in the past year (Table 2).

Table 2) Report on the relationship between the life expectancy questionnaire score and the demographic and occupational characteristics of individuals

	pa	tional characteristics			
Row	Variable	Status	Average score of the life expectancy questionnaire	Standard deviation	
1	Sex	Female Male	160.85 162.12	25.26 36.99	0.72
2	Marriago	Single	145.95	18.82	رم مرم 1 مرم
Z	Marriage	Married	168.34	33.62	<0.0001
3	Having children	Yes	170.47	38.20	<0.0001
3	maving ciniuren	No	152.32	19.09	<0.0001
4	Child living with the parent in question	Yes	170.47	38.20	< 0.0001
	omia nying with the parent in question	No	148.69	13.82	-0.0001
		General Practitioner	135.76	29.95	-
5	Educational field	Emergency Medicine	174.40	23.29	< 0.0001
3	Educational field	Specialist Specialist in Other		24.2	. \0.0001
		Fields	155.68	34.87	
6	Employment of the spouse in question	Yes	168.85	34.12	0.008
0	Employment of the spouse in question	No	148.35	7.36	0.008
		PhD	163.90	37.56	
7	Spouse's latest educational degree	Masters	172.08	9.74	0.56
		Bachelors	167.81	20.74	
8	Average income-activity ratio	Yes No	159.67	29.72	0.69
		Very little	161.76 149.14	31.90 36.71	
		A little	172.57	17.30	-
9	Satisfaction with one's work situation	Average	164.99	31.78	< 0.0001
,	badistaction with one 3 work situation	A lot	0	0	-
		Very much	0	0	
		Very little	135.14	38.90	-
		Little	155.98	17.40	<0.0001
10	Satisfaction with one's life situation	Average	166.71	21.88	. <0.0001
		A lot	213.22	15.08	-
		Very much	0	0	
		Very little Little	157.17	20.27 42.70	-
11	Desire to change career in current	Average	188.50 168.90	20.75	<0.0001
11	situation	A lot	171.55	21.16	
		Very much	145.13	36.81	
		Very little	145.01	38.98	
		Little	168.76	27.28	< 0.0001
12	Traveling to relieve job stress	Average	167.54	21.95	
	,	A lot	129.58	2.02	-0.0001
		Very much	191.00	0.00	<0.0001
		Very little	140.93	50.02	_
	Communicating with relatives to relieve	Little	172.56	26.20	-
13	job stress	Average	156.02	23.92	< 0.0001
	JOD 311 C33	A lot	162.16	22.52	-
		Very much	0	0	
		Very little	145.32	44.12	_
1.4	Francisco to relice to inlease	A little	165.38	21.38	<0.0001
14	Exercise to relieve job stress	Average	175.98	21.03	. <0.0001
		A lot	179.31 131.00	18.25 0.00	-
		Very much Not at all	117.15	42.44	
		Somewhat	152.78	25.06	•
15	Adequate sleep during the day and night	Rather	176.13	26.76	<0.0001
-	1 · · · · · · · · · · · · · · · · · · ·	Often	166.16	22.19	
		Completely	175.00	0	·
16	Specific physical or psychological illness	Yes	164.33	27.98	0.16
10		No	159.35	33.94	0.10
17	Specific illness of first-degree family	Yes	166.39	34.34	0.04
1/	members	No	158.74	29.66	0.04
18	Emergency shift status	Fixed shift	162.75	30.96	0.72
10	Emergency sum status	Rotating shift	161.16	31.78	0.72
		Morning	172.84	16.35	
		Evening	175.00	0	_
19	Emergency shift time	Night	130.57	43.07	<0.0001
17	Emergency sunt time	Combination of			\U.UUU1
		morning, evening and	164.78	28.19	
		night shifts	474.5	24.25	
20	Schedule of day off in the past year	Yes	161.65	31.95	0.55
20	Schedule of day off file bast veni				
20	Schedule of day off in the past year	No Loss than 4 nights	154.88 166.70	8.13	
21	Number of night shifts in the past year	No Less than 4 nights 5 to 8 nights	154.88 166.79 162.18	25.57 27.11	0.183

Toloei et al.

Stepwise multiple linear regression was used to test whether the variables of age, duration of marriage, number of children, total work experience, and work experience in the emergency department significantly predicted the life expectancy questionnaire score. The results of the fitted regression model test (model 1) were: Life expectancy questionnaire score = 4.84-294.91* (age) + 1.50* (marital duration) + 4.11* (number of children) + 2.83* (general work history) + 1.19* (emergency work history) The overall regression was statistically significant (p<0.0001, F (5.211) =11.64, R2 =0.216).

A stepwise regression was performed using backward elimination with a threshold of p<0.10. According to the regression results, the physicians' age, duration of marriage, general work history, and emergency work history significantly predict the life expectancy questionnaire score. Therefore, the mentioned variables were entered into the second regression model and the number of

children was excluded from this model. (Table 4) The regression results indicated that the age of the physicians significantly predicted the life expectancy questionnaire score (β = -5.04, p < 0.0001). While, the duration of the physicians' marriage significantly predicted the life expectancy questionnaire score (β = 1.79, p = 0.001). Also, the results of the analysis showed that the general work history of the physicians significantly predicted the life expectancy questionnaire score (β = 3.02, p < 0.0001). Finally, the work history in the emergency room of the physicians significantly predicted the life expectancy questionnaire score (β = 1.26, p = 0.01). (Table 3)

Below are the scatter plots (Figure 2) the P-P plot for the standardized regression residual (Figure 3) and the histogram plot (Figure 3). Pearson correlation was also used to examine the strength of the pairwise linear relationship between the above variables, the results of which are given in Table 4.

Table 3) Results of the stepwise linear regression model ((Backward) predictor variables for the total life expectancy score (Total HOL))

Model	Predictor variables entered	В	SE	Beta	T	p			
	Constant value	249.915	21.427	-	13.764	0.0001			
	Age	-4.849	0.747	-1.077	-6.493	0.0001			
	Duration of marriage	1.502	0.461	0.362	3.261	0.001			
1	Number of children	4.115	3.325	0.109	1.238	0.217			
	General work history	2.836	0.673	0.622	4.214	0.0001			
	Work history in the emergency department	1.195	0.461	0.215	2.594	0.01			
	NOTE: R=0.465, R2=0.216, ADJ.R2=0.198								
	Fixed amount	299.982	21.058	-	14.245	0.0001			
	Age	-5.045	0.731	-1.121	-6.904	0.0001			
	Duration of marriage	1.798	0.394	0.433	4.564	0.001			
2	Number of children	3.029	0.656	0.665	4.620	0.0001			
	Overall work history	1.267	0.457	0.228	2.771	0.006			
	Emergency work history	299.982	21.058	-	14.245	0.0001			

Table 4) Pearson correlation coefficient results

Variables	Life expectancy questionnaire score	Age	Duration of marriage	Number of children	Overall work history	Work history in the emergency department
Life Expectancy Questionnaire Score	1.000	-0.041	0.177	0.236	0.104	0.084
Age	-0.041	1.000	0.758	0.528	0.903	0.688
Duration of marriage	0.177	0.758	1.000	0.695	0.729	0.481
Number of children	0.236	0.528	0.695	1.000	0.578	0.400
General work history	0.104	0.903	0.729	0.578	1.000	0.596
Work history in the emergency department	0.084	0.668	0.481	0.400	0.596	1.000

8 Descriptive and analytical study of factors affecting life expectancy in doctors on duty in the emergency department




Figure 2) Scatter plot for the standardized regression residual for predicting the life expectancy questionnaire score.




Figure 3) Normal P-P plot for the standardized regression residual for predicting life expectancy questionnaire score




Figure 4) Standardized Regression Residual Histogram for Predicting Life Expectancy Questionnaire Score

JOURNAL OF POLICE MEDICINE

Vol.13, Issue 1, 2024

Toloei et al.

DISCUSSION

This study aimed to investigate the factors affecting life expectancy in shift physicians working in the emergency department.

The changes of recent decades have changed the work system and economy. Although shift work has been common among medical staff since the beginning, the shortage of experienced labor on the one hand and the economic needs of individuals on the other have made this issue more important in this group than ever before. In this study, unlike previous studies that were mainly focused on nurses, the factors affecting life expectancy and problems of shift work in general practitioners and specialists working in the emergency departments of public and private medical centers in Tehran were examined. As an organization providing medical services to citizens, medical centers should take measures to improve the quality and life expectancy and solve problems arising from the job position of clinical staff (medical staff), because having dynamic, motivated, hopeful, and purposeful staff, can better and faster achieve their long-term organizational goals and provide optimal services to their clients. Previous studies have shown that shift work has significant effects on sleep, depressed mood and anxiety, substance use, cognitive impairment, poor quality of life, and even suicidal thoughts, with clear sleep disorders often underlying the mental health consequences of shift work [20].

Asghari et al. (2013), in a study of 150 workers employed in an automotive industry in Tehran, showed that the highest prevalence of problems caused by shift work was related to disruption in the family, personal, and social lives of shift workers [21].

The findings of Mehdizadeh's study showed that nurses who had more night shifts in their work schedule and those who had less than 2 hours of sleep during night work, as well as nurses who did not participate in setting their work schedule and personnel working in ICU wards, face more hormonal changes [22]. In Kakui's study, a comparison of the temporal variation model of melatonin secretion in shift and fixed-duty nurses indicated that shift work is inherently unnatural and abnormal for humans, and by changing the pattern of melatonin secretion (a significant decrease in melatonin secretion at the peak, i.e. 4 a.m. and early morning, i.e. 7 a.m.), the sleep cycle, nighttime alertness, and daily activity will be disrupted [23].

In this study, the relationship between age, marital status, having children, whether the child lives with the parent in question, field of study, employment of the spouse in question, most recent educational

degree, proportion of average income to activity, level of satisfaction with one's work situation, level of satisfaction with one's life situation, desire to change careers in the current situation, traveling to relieve work stress, interacting with relatives to relieve work stress, exercising to relieve work stress, adequate sleep during the day and night, specific physical or psychological illness, specific illness of first-degree family members, day off shift in the past year, and number of night shifts in the past year were examined with life expectancy. Based on the results, the life expectancy test score was significantly related to marital status, parenthood, child living with the parent in question, field of study, spouse employment, child living with the parent in question, field of study, spouse employment, satisfaction with one's work situation, satisfaction with one's life situation, desire to change careers in the current situation, traveling to relieve job stress, interacting with relatives to relieve job stress, exercising to relieve job stress, adequate sleep during the day and night, specific illness of first-degree family members, and shift status in the emergency room. Also, the age of the physicians, duration of marriage of the physicians, general work history of the physicians, and work history of the physicians in the emergency room significantly predicted the life expectancy test score. The type of work shift had a significant relationship with life expectancy, meaning that people who mainly worked the morning or evening shift had higher life expectancy scores than people on rotating and night shifts. These results are consistent with the studies of Mirmohammadi et al. (2011), Ardakani et al. (2007), Ferri et al. (2016), and Barrau-Baumstarck et al. (2009) [24-27]. The above researchers believe that inappropriate shift work, due to its negative impact on social health, causes the individual to be deprived of regular and continuous social interactions due to the lack of a structured and predetermined life plan. Our study findings did not find a significant difference in the score of the life expectancy test of physicians in different groups based on the shift on a holiday in the past year and the number of night shifts. This result is in line with the findings of previous studies, including the works of Farahnaki et al. (2014), and Sabokroo et al. (2011), which indicate the negative impact of night shift work and official holidays on incompatible family life. Also, although the results of Farahnaki et al. (2014) and Asghari et al. (2013) point to the absence of shift workers at home on weekends and holidays and the poor quality of their time at home due to the inability to closely monitor household affairs due to fatigue from shift work, our study did not find a negative

Descriptive and analytical study of factors affecting life expectancy in doctors on duty in the emergency department

relationship with life expectancy in shift workers [21, 28, 29].

Barnes-Farrell (2010), and Barrau-Baumstarck et al. (2009) believe that shift workers should try to cope with and adapt to their job conditions; because if they do not adapt to the conditions, they will not be able to deal with problems that arise in personal, family, and professional areas, and this will cause burnout for the shift worker [27,30]. The study had limitations, including the breadth and multifactorial nature of the factors affecting life expectancy, which could potentially have affected the results of the study. Future studies should examine the impact of variables such as employment status and diet on life expectancy. Another limitation of this study was the data collection, which may have biased the responses of respondents through self-reporting.

CONCLUSION

The results of this study revealed that in appropriate shift work can disrupt the personal, family, and professional lives of medical personnel. In addition, various factors, including demographic characteristics, shift work, income, sleep, living conditions, travel, exercise, contact with relatives, specific physical or psychological illnesses, and specific illnesses of family members, affect the life expectancy of shift-working physicians. It seems that reducing the working hours of personnel, avoiding fixed night shifts or excessive shifts on public holidays among physicians working in the emergency department can prevent burnout and reduce life expectancy. In this case, the social capital of personnel will increase and job satisfaction and the quality of patient care will also increase. Considering the sensitive nature of the medical profession, especially physicians working in the emergency department, and the need for a 24-hour presence in the country's medical sector, it is necessary to consider measures to maintain the health of this group of physicians, including alternative incentives.

Clinical & Practical Tips in POLICE MEDICINE:

Authors' Contribution: All authors participated in the initial writing and revision of the article, and all accept responsibility for the accuracy and completeness of the content of the article with final approval.

Conflict of interest: The authors stated that there is no conflict of interest in the present study.

Financial Sources: There was no sponsorship in this article.



نشریه طب انتظامی



مسترسی آزاد 👌



مقاله اصيل

بررسی توصیفی تحلیلی عوامل مؤثر بر امید به زندگی در پزشکان نوبت کار بخش اورژانس

- ٔ گروه روانشناسی تربیتی، دانشکده علوم انسانی، دانشگاه آزاد اسلامی واحد اراک، اراک ایران.
- مرکز تحقیقات تروما در عملیات پلیس، معاونت بهداشت، امداد و درمان فراجا، تهران، ایران.
- گروه حقوق، دانشکده حقوق، علوم سیاسی و زبانهای خارجی، دانشگاه آزاد اسلامی واحد مشهد، مشهد، ایران.

حكىدە

اهداف: نوبت کاری در قالبها و یا نظامهای گوناگونی که هر یک دارای ویژگیها، مزایا و معایب خاص خود میباشند اجرا میشود و میتواند اثرات نامطلوبی بر جنبههای مختلف زندگی انسان بجا گذارد. کاهش امید به زندگی بر سلامت جسمی و روحی اثرگذار بوده و به عنوان پیامد نهایی استرس شغلی، در بسیاری از مشاغل نوبت کار، از جمله پرستاری مشاهده میشود. هدف از این مطالعه، بررسی عوامل مؤثر بر امید به زندگی، در پزشکان نوبت کار بخش اورژانس است.

مواد و روشها: این مطالعه از نوع مطالعات توصیفی- تحلیلی و مقطعی بود که در سال ۱۴۰۲ در مرکز تحقیقات کاربردی معاونت بهداد فراجا انجام شد. جمعیت مورد مطالعه، یزشکان شیفتی بخش اورژانس ۵۰ مرکز بیمارستانی و درمانگاهی دانشگاههای علوم یزشکی ایران و البرز، با حداقل ۶ ماه سابقه فعالیت شغلی پیوسته، با شیفت ثابت و در گردش، در سال ۱۴۰۱ بودند. نمونهگیری به روش طبقهای، با تعداد نمونهٔ ۳۰۰ نفر (سطح اطمینان ۹۵ درصد) صورت گرفت. بدین منظور، در شیفتهای مختلف کاری مراجعه و ضمن توضیح در خصوص اهداف مطالعه و آموزش دستورالعمل، پرسشنامهٔ امیدواری میلر و نحوهٔ تکمیل آن، جهت جمعآوری اطلاعات دموگرافیک، در اختیار پزشکان قرار گرفت. سیس دادهها، با استفاده از نرمافزار SPSS 26 تحلیل شد. جهت مقایسهٔ دادهها، از تستهای مناسب پارامتریک و غیرپارامتریک، از جمله t-test و ANOVA استفاده شد. همچنین، ارتباط بین متغیرهای کمی با استفاده از همبستگی پیرسون و رگرسیون خطی بررسی شد.

یافتهها: ۳۱۳ نفر (۱۵۸ زن) شامل ۹۴ پزشک عمومی (۳۰ درصد)، ۱۹۷ پزشک متخصص اورژانس (۶۲/۹ درصد) و ۲۲ پزشک متخصص در سایر رشتهها (۷ درصد)، با میانگین سنی ۷/۴۷ ± ۴۴/۰۷ وارد مطالعه شدند. میانگین نمرهٔ آزمون امید به زندگی، اختلاف معناداری در زيرگروههاي مختلف وضعيت تاهل (p<-/٠٥١)، فرزندداري (p<-/٠٥١)، زندگي كردن فرزند با والد مورد سؤال (p<-/٠٥١)، رشتهٔ تحصيلي (٥٠٠١-)، شاغل بودن همسر (۸۰۰/هٔ p)، میزان رضایت از وضعیت کاری خود (p۰۰/۰۰۱)، میزان رضایت از وضعیت زندگی خود (p۰/۰۰۱)، تمایل به تغییر حرفه در وضعیت فعلی (p<۰/۰۰۱)، مسافرت برای رهایی از استرس شغلی (p<۰/۰۰۱)، مراوده با نزدیکان برای رهایی از استرس شغلی (p<-/٠٠١)، ورزش برای رهایی از استرس شغلی (٥٠٠/٠٠١)، خواب مناسب در طی شبانه روز (p<-/٠٠١)، بیماری خاص اعضای درجه اول خانواده (p=-/۰۴) و وضعیت شیفت در اورژانس (p<-/-/0) داشت. همچنین، نتایج رگرسیون خطی نشان داد که سن پزشکان، مدت زمان تأهل پزشکان، سابقه کار کلی پزشکان و سابقه کار در اورژانس پزشکان، به طور قابل توجهی نمره آزمون امید به زندگی را پیش بینی کرد (۲۱۶-۳2 و و .(p<°/>/°°1 9 F(Δ,Υ11)=11/۶۴

نتیجهگیری: شیفت کاری نامناسب، موجب کاهش امید به زندگی و به دنبالش، اختلال در زندگی فردی، خانوادگی و شغلی پرسنل کادر درمان می شود. در ضمن، ویژگیهای جمعیت شناختی، شیفت کاری، خواب مناسب شبانه، وضعیت زندگی، مسافرت، ورزش، مراوده با نزدیکان، بیماری خاص جسمی یا روانشناختی و بیماری خاص اعضای خانواده، بر امید به زندگی اثر معنیداری دارد. همچنین افزایش شیفت کاری پزشکان شاغل در بخش اورژانس، به ویژه شیفتهای شب، موجب کاهش معنیدار امید به زندگی این گروه از پزشکان میشود.

كليدواژهها: برنامه نوبت كارى، روانسنجى، رضايت شخصى، اميد، اورژانس بيمارستان

تاريخچه مقاله:

دریافت: ۱۴۰۲/۰۲/۱۵

یذیرش: ۱۴۰۳/۰۴/۲۹ انتشار: ۱۴۰۳/۰۵/۰۳

نويسنده مسئول *:

آدرس: تهران، خیابان ولیعصر(عج)،بالاتر از میدان ونک، روبروی ظفر، بیمارستان حضرت ولیعصر(عج)،طبقه ینجم، مرکزتحقیقات کاربردی معاونت بهداد فراجا، کد پستی:

بست الكترونيكي: hamedaghdam@gmail.com

نحوه استناد به مقاله:

Toloei S, Aghdam H, Shirzad O, Vakilipour P. Descriptive and analytical study of factors affecting life expectancy in doctors on duty in the emergency department. J Police Med. 2024;13:e13.

بقدمه

نوبت کاری (Shift Working) از جمله پدیدههای اجتماعی است که ریشه در تاریخ انسان دارد. امروزه به دلایل اقتصادی و تکنولوژیک، سیستمهای نوبت کاری، به صورت جزئی از جامعهٔ شبانهروزی، در سراسر جهان درآمده است. چراکه، بایستی به نحوی تقسیم کار صورت گیرد، که در همه ساعات شبانه روز، افراد فعالی در محل کار حضور داشته باشند و در بسیاری از صنایع و کارخانجات صنعتی-تولیدی، همچنین در برخی مشاغل خدماتی مانند پزشکی، پرســتارى، آتشنشــانى، انتظامــى، خدمــات آب، بــرق، تلفــن و... اجـرا میشـود [۱]. قالبهـا یـا نظامهـای گوناگونـی از نوبت کاری وجود دارد که هر یک، دارای ویژگیها، مزایا و معایب خاص خود هستند. دو نظام کلی نوبت کاری عبارتند از، نظام نوبت کاری ۱۲ ساعته (دونوبتی) و نظام ۸ ساعته (سه نوبتی). یک تعریف کلی از نوبت کاری، عبارت است از الگوی غیرمعمول کاری در مقایسه با روزکارها، که این تعریف شامل چند نوع نوبت کاری میشود، ازجمله شیفتهای ثابت که در این نوع، عدهای به طور ثابت در نوبت صبح، عصر یا شب مشغول فعالیت میباشند. شیفتهای چرخشی، که افراد به طور چرخشی در شیفتهای صبح، عصر و شب مشغول به کار هستند و شامل شیفتهای چرخشی رو به جلو (جهت عقربههای ساعت) و شیفت چرخشی رو به عقب، که عکس حالت اول انجام میگیرد و میتواند به صورت چرخش سریع (۲ تـا ٣ روز)، چرخـش متوسـط (١ هفتـه) و چرخـش كنـد (٢ تـا ۳ هفتـه) بـرای هـر شـیفت کاری در نظـر گرفتـه شـود [۲]. بـه دلیل اینکه افراد نوبتکار مجبورند هم در روز، که جامعه فعـال اسـت و هـم در شـب، كـه جامعـه در خـواب بـه سـر مىبرد، بيدار بوده و به فعاليت مشغول باشند، اين پديده، امروزه به عنوان یک عامل تهدیدکنندهٔ سلامت تلقی شده که میتواند اثرات نامطلوبی بر جنبههای گوناگون زندگی انسان بجا گذارد [۳].

مطالعات متعددی با هدف بررسی اثرات منفی نوبت کاری بر عملکرد، سلامت و کیفیت زندگی کارگران نوبت کار انجام شده است که برخی از این مطالعات ارتباط نوبت کاری و اختلال خواب، مشکلات معدی-رودهای، بیماریهای قلبی- عروقی و اختلالات روحی- روانی را نشان دادهاند [۴].

نوبت کاری در حرفه پزشکی، در مؤسسات و مراکز بهداشتی- درمانی و بهویژه بخش اورژانس، دارای پیشینهای دیرین بوده و از دیرباز، به صورت یک امر شایع درآمده است. از آنجایی که بیماریهای حاد، بیثباتی وضعیت جسمانی و روانی و وابستگی بیماران به دیگران، باعث الزام ارائه خدمات پیوسته و مداوم میگردد، بنابرایین ذات چنین خدماتی، به دور از برنامهریزی قبلی است و لازم است و افرادی به صورت مستمر و در طول شبانهروز، برای ارائه خدمات حضور داشته باشند. بر پایه مطالعات انجام گرفته، خدمات حضور داشته باشند. بر پایه مطالعات انجام گرفته،

شیوع نوبت کاری در کشورهای اروپایی ۱۵ تا ۲۰ درصد و در آمریکا ۲۰ درصد نیبروی کار است [۵]. ایبن میزان شیوع، دلیل مناسبی برای مطالعه و بررسی اثرات منفی بر سلامتی، ایمنی و کارایی کارگران خواهد بود. نوبت کاری یکی از قدیمی تریب چالشهای کارکنان مراقبتهای بهداشتی است و می تواند اثرات سوء فیزیولوژیک، روانی و اجتماعی را بر افراد نوبت کار برجا گذارد [۶].

پیامدهای اجتماعی، روانی و خانوادگی شامل تحریکپذیری، افسردگی، استرس و اختالال در روابط اجتماعی، از جمله پدیدههای روحی و روانی ناشی از تداخل شیفت کاری با زندگی اجتماعی و خانوادگی افراد است. شیفت کاری میتواند بر روی زندگی خانوادگی افراد اثر منفی داشته باشد و زندگی اجتماعی آنها را مختل سازد، بویژه، وقتی برنامهٔ شیفت کاری بهگونهای است که فرد در اواخر بعد از ظهر و شب، از خانه دور باشد [۶].

اگرچه در کشورهای پیشرفته، مطالعات متعددی در زمینهٔ اثرات ساعات کار زیاد در نوبتهای ۱۲ ساعته و یک هفتهای در گروههای شغلی مختلف و همچنین، پیامدهای کار در نوبتهای ۱۲ ساعته و دورههای ۲ هفتهای انجام شده و در آنها اثرات ساعات کار طولانی و برنامهٔ نوبت کاری بر روی عملکرد، سلامتی، کیفیت زندگی و حوادث شغلی كنكاش شده است [٧]، اما نتايج اين تحقيقات به علت تفاوتهای سازمانی، محیطی و فرهنگی، نمیتواند الگوی کاربردی کاملاً مناسبی برای کشورمان در اختیار گذارد. همچنین، براساس مطالعات انجام شده بر روی برنامههای نوبت کاری، مشخص شدہ کہ یک برنامہ کاری با طراحی ضعیف میتواند بر کیفیت مراقبتهای بهداشتی، رضایت بیمار، طول دوره بستری، غیبت از کار، اثر بخشی هزینهها و بهرهوری تأثیر گذارد [۸][. اکثر این مطالعهها، بر روی کادر درمانی پرستاری و سایر افراد شاغل در بخشهای درمانی انجام شده است و بهندرت این مطالعه، بر روی پزشکان انجام شده است. در سالهای گذشته، با گسترش بیش از قبل رشته تخصصی طب اورژانس و الزام مراکز درمانی به حضور متخصص طب اورژانس در بخش اورژانس، بررسی این موضوع ضرورت بیشتری پیدا کرده است. در مراکز درمانی درمانگاهی و بخشهای بیمارستانی، پزشکان عمومی نیز حضور داشتهاند که بررسی تأثیر شیفت کاری بر روی این قشر نیز ضروری است.

امید و امیدواری به زندگی، از مفاهیمی هستند که با تطابق، ایمان و توانمندی مرتبط بوده، از طریق که با تطابق، ایمان و توانمندی مرتبط بوده، از طریق تحمل پذیرکردن شرایط موجود میتواند به عنوان پاسخی مؤثر به عوامل تنشزا در نظر گرفته شود. امیدواری از طریق تجارب موفقیت آمیز زندگی افزایش می یابد و به واسطه تجارب شکست تقلیل می یابد. کسانی که سبک تبیین مثبت دارند، شادی، امید و رضایت بیشتری از زندگی دارند [۹]. امید به زندگی، به عنوان یک عامل قدرت بخش، چند بعدی، پویا و قدرتمند در سازگاری با مشکلات و حتی

بررسی توصیفی تحلیلی عوامل مؤثر بر امید به زندگی در پزشکان نوبتکار بخش اورژانس

ناسازگازیها مؤثراست. امید به زندگی، یکی از مفاهیم اساسی روان شناسی مثبت نگر و عاملی انرژی بخش در زندگی افراد است. افراد با امید به زندگی بالاتر، قدرت بیشتری برای نیل به اهداف خود دارند و هنگامیکه با مانعی برخورد می کنند، می توانند انگیزه خود را حفظ و از راههای جانشین استفاده کنند؛ اما افراد ناامید چون راههای کمی دارند، در رویارویی با موانع انگیزه خود را از دست میدهند. همچنین امید، عاملی به عنوان انرژی روانی متمرکز بر هدف و احساس مصممبودن در دستیابی به هدف تعریف شده است. افراد دارای امید راهبردی بالا، به منظور مقابله با موانع احتمالی، راههای متعددی را در نظر می گیرند. در مقابل، افراد با امید راهبردی پایین، احتمال پیداکردن راه حلی ممکن، برای رسیدن به اهدافشان را ندارند. افرادی که میتوانند تعارضات موجود در رابطه را با به کارگیری روشهای مثبت و استفاده کمتر از تعاملات منفی مدیریت کنند، فضایی ایجاد میکنند که در آن فرصت بیشتری برای خودافشایی و توافق در مورد مشکلات زندگی وجود خواهد داشت [۱۰].

Schneider و همـکاران، امیـد را سـازهای دوبُعـدی دانستهاند: یکی توانایی طراحی راههایی به سوی هدفهای مطلوب با وجود موانع (تفکر راهبردی) و دیگری توانایی ایجاد انگیزه (تفکر عاملی) برای شروع و تداوم حرکت در این گذرگاهها [۱۱]. آنها امید را به عنوان مجموعهای شناختی توصیف نمودند که مبتنی بر احساس موفقیت، ناشی از منابع گوناگون (تصمیمهای هدفدار) و مسیرها (شیوهٔ انتخاب شده برای رسیدن به اهداف) است. افرادی کے دارای امیے بالایے هستند، دارای یے حس مدیریت تعییان هدف، هماراه با توانایی بارای تولید برنامههایی برای رسیدن به اهداف خود هستند. Schneider و Michael همچنین مطرح کردند که امید ممکن است یک اثـر تعدیل کننـده داشـته باشـد. بـه طـوری کـه، افـرادی کـه دارای امیدواری بالا هستند، نسبت به آنهایی که امیدواری کمتری دارند، توانایی بیشتری برای پیادهسازی استراتژی سازگاری و مقابلهای با بحرانها دارند. آنهایی که امید بیشتری دارند، قادر به مفهومسازی اهداف خود بهروشنی بوده و احساس اعتماد به نفس و قابلیت بیشتری برای مدیریت مؤثر بر عوامل استرسزا دارند و به خوبی میدانند که برای مقابله با هر استرسی، باید برنامه خاصی داشته

امید به زندگی، یکی از متغیرهای اساسی است که میتواند در تابآوری شرایط ویشهٔ کاری کادر درمانی، سازگاری با شرایط استرسزای محیط اورژانس و تبعیت از نظامهای علمی و اداری کمک کند. امید، از مهمترین مباحث در روان شناسی مثبت است. روان شناسان مثبت نگر معتقدند، امید میتواند از افراد در برابر رویدادهای تنـشزای زندگـی حمایـت کنـد. امیـد شـامل ادراکات شـخص دربارهٔ تواناییهایـش بـرای تعییـن اهـداف روشـن، تدویـن

طرحهایی برای رسیدن به این اهداف، یافتن و حفظ انرژی و انگیزش مورد نیاز برای رفتن به دنبال آن اهداف و پیگیری آنها است. امید شامل تصورات فرد به آینده مثبت است و موجب افزایش تلاش وی برای بهبود شرایط می شود. شواهدی وجود دارنید که نشان می دهند، بین بالا بودن مینزان امیند افتراد با موفقیت آنها در فعالیتهای مختلف، کیفیت زندگی و سلامت جسمی و روانی بهتر، ارتباط وجود دارد [۱۳]. امید در مواجهه مؤثر با مشکلات و استرس ناشی از آن، نقش مهمی دارد و به عنوان عاملی بالقوه نیرومند در بهبود و سازگاری بیماران مؤثر است. از همین رو، امید با افزایش هدفمندی در زندگی شخصی و کاری همـراه اسـت [۱۴]. در مطالعـهای Cao و همـکاران [۱۵]، نشان دادهاند در کادر درمانی، برخورداری از امید بالاتر پیشبینی کننده مشارکت بهتر آنها در فرآیند درمان است. همچنین، Clayton و همکاران نیز، در یک مطالعه متاآنالیز نشان دادنید کیه بیه طبور کلی، حیس امییدواری، فاکتبوری مهم برای انگیزهداشتن در فرآیند درمان است و در افرادی کے امید بے زندگی بیشتری دارند، رضایتمندی بهتر

با توجه به مطالعات قبلی، میتوان دریافت که نوبتکاری، با ایجاد مشکلات روحی و روانی در افراد، میتواند باعث تضعیف امیدواری آنها شود که این موضوع در قشر پزشکان، میتواند آثار زیانباری را بر بیماران و سیستم درمانی کشور داشته باشد. بنابراین، این مطالعه با هدف بررسی عوامل مؤثر بر امید به زندگی در پزشکان شیفتکار شاغل در بخش اورژانس انجام شد.

روش کار

این مطالعه از نوع توصیفی- تحلیلی بود. جمعیت مورد مطالعـه، پزشـکان نوبـت کار شـاغل در ۵۰ مرکـز بیمارسـتانی و درمانگاهـی مربـوط بـه دانشـگاههای علـوم پزشـکی ایـران و البرز بود كه در كل شامل ۱۰۰۰ نفر بودند. افراد مورد مطالعه، از بین پزشکان دارای حداقل ۶ ماه سابقه فعالیت شغلی پیوسته، با شیفت ثابت و در گردش در سال ۱۴۰۱ انتخاب شدند. با توجه به بررسیهای انجام شده، تعداد یزشکان مشغول در اورژانس مراکز مورد مطالعه مشابه بودنـد. بنابرایـن، از روش نمونه گیـری تصادفـی سـاده اسـتفاده شد. برای این منظور، ابتدا فهرستی از تمام پزشکان شاغل در اورژانس در همه مراکز تهیه شد. سیس، به هر پزشک کد اختصاصی برای انتخاب تصادفی اختصاص یافت. در نهایت، با استفاده از نرمافزار اکسل، به تعداد مورد نیاز، کدهای تصادفی تعیین شد و نمونههای مطالعه انتخاب شدند. حجم نمونه مورد نیاز برای مطالعه، با در نظر گرفتن سطح اطمینان ۹۵ درصد و مقدار z برابر با ۱/۹۶، خطای نمونهگیری مجاز اصلاحشدهٔ ۵/۸ درصد، پیشبینی نسبت موفقیت (p) برابر با ۵/۰ و با استفاده از فرمول برابر بــا ۲۸۵ نفـر محاســبه شــد و بــا $n=Z2 \times p \times (p-1)/E2$

در نظر گرفتن احتمال ریزش آماری، ۳۰۰ نفر در نظر گرفته شد. همچنین، به علت بار کاری زیاد پزشکان، و به منظور جلوگیری از ریزشهای احتمالی در پاسخگویی افراد، تعداد بیشتری (۳۵۰ مورد) پرسشنامه توزیع گردید.

معیار ورود به مطالعه وضعیت شغلی (پزشکان دارای حداقل ۶ ماه سابقه فعالیت شغلی پیوسته در شیفتهای ثابت و در گردش)، نوع شغل (فقط پزشکان شاغل در اورژانس مراکز بیمارستانی و درمانگاهی مشخص شده به عنوان جامعه مورد مطالعه) و اشتغال به کار (پزشکان در حال فعالیت و اشتغال به کار در زمان نمونهگیری) بود. از طرفی، عدم تکمیل پرسشنامه، مرخصی یا غیبت پزشکان، عدم تجربه یا تخصص مرتبط و مشکلات سلامتی، معیار خروج در نظر گرفته شد.

پرسشنامهای شامل دو بخش اطلاعات دموگرافیک و پرسشنامه امید به زندگی (HOL) به عنوان ابزار جمعآوری دادهها استفاده شد. پرسشهای ویژگیهای دموگرافیک و خصوصیات فردی شامل جنسیت، سن، سابقهٔ کار، سابقهٔ فعالیت در بخش اورژانس، شیفت کاری، درآمد، خواب، وضعیت زندگی، مسافرت، ورزش، مراوده با نزدیکان، بیماری خاص خود و اعضای خانواده در گروههای شغلی بیماری خاص خود و اعضای خانواده در گروههای شغلی مختلف پزشکان در بخش اورژانس بود.

جهت سنجش ميزان اميدواري افراد مورد مطالعه، از پرسشنامهٔ امید میلریا پرسشنامهٔ امیدواری میلر (MHS) استفاده شد. این پرسشنامه، رایجترین ابزار اندازهگیری امید به زندگی است که از ۴۸ ماده جداگانه در مورد احساسها و نگرشها تشکیل شده و جنبههای مختلف امید به زندگی را میسنجد. این پرسشنامه ترجمهشده، از روی یک ابزار به زبان انگلیسی است که تاکنون در ایران اجرا نشده است و نیازمند اعتباریابی (سنجش روایی و پایایی) است. Miller روایی این پرسشنامه را مطلوب گـزارش کـرده و پایایـی آن را طبـق آلفـای کرونبـاخ بـالای ۸۰ ذکر کرده است [۱۷]. پاسخهای این پرسشنامه، در یک پیوستار ینج درجهای (بسیار مخالف= ۱، مخالف= ۲، بیتفاوت= ۳، موافـق= ۴، بسـیار موافـق= ۵) تنظیـم شـده اسـت. در آزمـون میلر، دامنه امتیازات کسب شده از ۴۸ تا ۲۴۰ متغیر است. ۱۲ جمله از پرسشامه میلر از مادههای منفی تشکیل شده که این شمارهها، در ارزشیابی و نمرهگذاری برعکس نمره میگیرنــد [۱۷].

آزماییش مقدماتی ایین معیار، روی ۷۵ نفراز دانشجویان در آمریکا اجرا شد. این آزمایش، مادههای امید را به طور روشن از نظر سطح آمادگی، ویژگی اعتبار تحت بررسی قرار داد. ضریب پایایی ۹۵ درصد و اعتبار (تست مجدد با فاصلهٔ دو هفته) ۸۷ درصد و میانگین امیدواری ۱۷۵/۶ بر آورده شده است [۱۸]. این آزمون برای اولین بار در ایران، توسط سمیعی (۱۳۶۸) بر روی ۳۰ نفر از دانشجویان پسر رشته پرستاری اجرا شد و به منظور تعیین پایایی آن، از روش آزمون مجدد، با فاصلهٔ یک هفته استفاده شد.

ضریب همبستگی پیرسون بین دوسری از نمرات آزمون ۷۵ درصد برآورده شده است و میانگین آن ۱۵۱/۳ است. به منظور اطمینان از پایایی آزمون، این پرسشنامه بر روی تعداد بیشتری (۱۵۲ نفر) از دانشجویان دختر و پسر مراکز تربیت معلم تهران اجرا گردید که ضریب آلفای کرونباخ ۱۹۲۸ درصد به دست آمده ۱۹۶ درصد به دست آمده و میانگین به دست آمده ۱۹۶ بود [۱۹].

پس از اخذ مجوز، جهت تکمیل پرسشنامهها و جمعآوری دادهها، در شیفتهای مختلف کاری به درمانگاه و بیمارستان مراجعه و ضمن توضیح در خصوص اهداف مطالعه و آموزش دستورالعمل و نحوهٔ تکمیل، پرسشنامه به صورت آنلاین در اختیار پزشکانی که همکاری خود را اعلام کرده بودند، قرار گرفت. تمامی سؤالات پرسشنامه، به صورت خودگزارش، توسط افراد تکمیل شد. در این مطالعه، ابتدا میزان امید به زندگی مشخص گردید و سپس، میانگین امتیاز امید به زندگی در بین پزشکان و شیفتهای کاری مختلف به دست آمد، اطلاعات دموگرافیک افراد و سؤالات مربوط به علل احتمالی تأثیرگذار بر امید به زندگی، مورد تجزیه و تحلیل احتمالی تأثیرگذار بر امید به زندگی، مورد تجزیه و تحلیل قرار گرفت.

ملاحظات اخلاقی. این مطالعه، مطابق با اصول اخلاقی مندرج در اعلامیه هلسینکی انجام شد. تأییدیه اخلاقی از کمیتهٔ اخلاق در پژوهشهای زیست پزشکی معاونت بهداد فراجا با کد اخلاق زیست پزشکی معاونت بهداد فراجا با کد اخلاق شد. همهٔ مسرکتکنندگان، قبل از ورود به مطالعه، رضایت آگاهانه کتبی ارائه دادند. شرکتکنندگان، از حق خود برای انصراف از مطالعه، در هر زمان بدون هیچگونه پیامدی مطمئن شدند. تمام دادههای جمعآوریشده، برای محافظت از محرمانهبودن شرکتکنندگان، ناشناس شدند.

تجزیه و تحلیل دادهها. نرمالبودن توزیع دادههای کمی با استفاده از تست کولموگروف اسمیرنوف بررسی شد. جهت مقایسهٔ دادههای با توزیع نرمال، از تستهای پارامتریک از جمله t-test و ANOVA و جهت مقایسهٔ دادههای با توزیع غیرنرمال، از تستهای غیرپارامتریک استفاده شد. برای بررسی ارتباط بین متغیرهای کمی از همبستگی پیرسون و رگرسیون خطی T گام به گام، استفاده شد. در تمام آنالیزهای آماری، سطح value وکمتر از ۵۰/۰ شد. تمام تحلیلهای آماری با استفاده از نرمافزار SPSS 26 انجام شد.

بافتهها

با توجه به معیارهای ورود به مطالعه ۳۱۳ نفر پزشک شامل ۹۴ پزشک متخصص اورژانس (۶۲/۹ درصد) ۱۹۷ پزشک متخصص اورژانس (۶۲/۹ درصد) و ۲۲ نفر پزشک متخصص در سایر رشتهها (۷/۰ درصد) وارد مطالعه شدند که ۱۵۸ نفر زن ۵۰/۵۰ درصد) و مدا نفرمرد (۴۹/۵۰ درصد) با میانگین سنی

وره ۱۳، شماره ۱، ۱۴۰۳ نشریه علمی پژوهشی طب انتظامی

بررسی توصیفی تحلیلی عوامل مؤثر بر امید به زندگی در پزشکان نوبتکار بخش اورژانس

٧٠/۵	۱۷۰	دكترا		
۵/۰	۱۲	فوق ليسانس	۔ آخرین مدرک تحصیلی همسر	Υ
۲۴/۵	۵۹	ليسانس	-	
۴۲/	146	بلی	بیماری خاص جسمی یا	Λ
۵۷/۲	۱۷۹	خير	روانشناختی	^
۳۵/۸	117	بلی	بیماری خاص اعضای درجهٔ اول	۹
54/4	۲۰۱	خير	خانواده	`
14/4	۴۲	بلی	الما الله على الما الما الما	١.
ለ۶/۶	271	خير	تناسب میانگین درآمد با فعالیت -	10
۲۰/۱	۶۳	شيفت ثابت		.,,
٧٩/٩	۲۵۰	شیفت در گردش	نوع نوبت کاری در اورژانس	11
۶/۱	19	صبح		
۰/۳	١	عصر	_	
11/٢	٣۵	شب	زمان نوبت کاری در اورژانس	14
۸۲/۴	۲۵۸	ترکیبی شیفت صبح، عصر		
	1ωλ	و شب		
97/4	۳۰۵	بلی	شیفت روز تعطیل در یک سال	۱۳
۲/۶	٨	خير	گذشته	
18/8	۵۲	کمتر از ۴ شب		
۵۵/۹	۱۷۵	۵ تا ۸ شب	تعداد شیفت شب در یک سال ⁻ گذشته	16
۲۷/۵	٨۶	ىىشتر از ۸ شب	ددسته	


همچنین، در بررسی میـزان رضایـت از زندگـی، ۳/۳ درصد از افراد رضایت زیاد داشته و میزان رضایت از کار، ۳۸/۳ درصـد بسـیار کـم، ۳۳/۹ درصـد کـم، و ۲۷/۸ درصـد متوسط بود. هیچ یک از شرکت کنندگان در این مطالعه، رضایت زیاد یا بسیار زیاد از کار خود نداشتند. در بررسی تمایل افراد به تغییر شغل، ۳۴/۲ درصد تمایل بسیار زیاد، ۳۳/۵ درصـد تمایـل زیـاد، ۱۵/۳ درصـد تمایـل متوسـط، ۵/۸ درصد تمایل کم و ۱۱/۲ درصد تمایل بسیار کم به تغییر شـغل داشـتند. در بررسـی عوامـل رهایـی از اسـترس شـغلی، تعداد اندکی از افراد جهت کاهش استرس شغلی، به صورت زیاد و بسیار زیاد تمایل به مسافرت، ورزش و وقت گذرانی با نزدیکان داشتند (به ترتیب ۶/۷ درصد، ۶/۴ درصد و ۱۴/۱ درصد). در بررسی خواب شبانه، حدود ۲۳/۶ درصد از افراد خواب مناسب را در طول شب دربیشتر مواقع گزارش کردند و ۸/۶ درصد از افراد خواب شبانه مناسب اصلاً نداشتند. جزئیات نتایج پرسشنامه در قالب نمودار ستونی لیکرت در شکل ۱ آورده شده است.

۴۴/۰۷±۷/۴۷ سال بودند که از این بین ۶۹/۳ درصد متأهل (۳۰/۷ درصد مجرد) بود. همچنین ۵۰/۵ درصد از افراد شرکت کننده در مطالعه صاحب فرزند بودند که از بین افراد صاحب فرزند، ۷۶/۳ درصد با فرزندشان زندگی می کردند. ۹۱/۳ درصد از افراد مورد مطالعه همسر شاغل داشتند.

نتایج بررسی وضعیت سلامتی روشن ساخت که درصد از افراد مورد مطالعه از بیماری جسمی یا روانی ۴۲/۸ درصد از افراد سابقهٔ بیماری میردند. همچنین ۳۵/۸ درصد از افراد سابقهٔ بیماری خاص را در نزدیکان درجه یک خود گزارش کردند. در بررسی نوع نوبت کاری، بیشتر افراد به صورت شیفتهای گردشی مشغول به کار بودند (۲۹/۹ درصد). شیفتهای ثابت شب در بین انواع شیفتهای ثابت بیشترین فراوانی را داشت کمتر از ۴ شیفت در ماه ۱۶/۶ درصد کمتر از ۴ شیفت در ماه ۱۶/۶ درصد در ماه ۱۶/۶ درصد بین ۴ تا ۸ شیفت در ماه، و ۵/۷۸ درصد بیش از ۸ شیفت در یک ماه داشتند. می موثر افراد در یک سال گذشته، شیفت روز تعطیل داشتند. در بررسی تناسب حقوق با کار، ۸۶/۶ درصد از افراد معتقد به عدم تناسب حقوق با کارشان درصد در جدول ۱ نشان داده شده است.

جدول ۱) مشخصات جمعیت شناختی و وضعیت شغلی افراد مورد مطالعه

درصد	تعداد	وضعيت	متغير	ردیف
۵۰/۵	۱۵۸	زن		
۴۹/۵	۱۵۵	مرد	جنسیت	١
۳۰/۰	916	پزشک عمومی		
84/9	197	متخصص طب اورژانس	رشتهٔ تحصیلی	۲
٧/٠	44	متخصص در رشتههای دیگر	رسته تحصیتی	1
۶۹/۳	717	متاهل	ı fı	٣
۳۰/۷	9,5	مجرد	وضعيت تأهل	Γ.
۵۰/۵	۱۵۸	بلی		۴
۴٩/۵	۱۵۵	خير	صاحب فرزند	۲
٧۶/٣	۱۵۸	بلی	زندگی کردن فرزند با والد مورد	Δ
۲۳/۷	k 9	خير	سؤال	ω
91/19	۲۰۹	بلی		۶
A/Y	۲۰	خير	شاغل بودن همسر	



شكل ۱) نمودار ليكرت نتايج پرسشنامهٔ اخذشده از افراد مورد مطالعه

نتاییج تست Kolmogorov Smirnov نشان داد که دادههای متغیرهای سن، مدت زمان تأهل، سابقه کلی کار، سابقه کار در اورژانس، و نمرهٔ پرسشنامهٔ امید به زندگی، توزیع نرمال دارند. بنابراین، جهت مقایسهٔ میانگین نمرهٔ پرسشنامهٔ امید به زندگی، بین گروههای مختلف از تست Sample t-test یا ANOVA استفاده شد. نتایج نشان داد که میانگین نمرهٔ پرسشنامهٔ امید به زندگی در زیرگروههای مختلف وضعیت تأهل، فرزندداری، زندگی فرزند با والد مورد سؤال، رشتهٔ تحصیلی، شاغلبودن همسر، زندگی کردن فرزند با والد مورد سؤال، میزان رضایت از وضعیت کاری خود، میزان

وضعیت فعلی، مسافرت برای رهایی از استرس شغلی، ورزش مراوده با نزدیکان برای رهایی از استرس شغلی، ورزش برای رهایی از استرس شغلی، ورزش برای رهایی از استرس شغلی، خواب مناسب در طی شبانه روز، بیماری خاص اعضای درجه اول خانواده و وضعیت شیفت در اورژانس، اختلاف معناداری دارد. ولی میانگین نمرهٔ پرسشنامهٔ امید به زندگی، در زیرگروههای مختلف جنسیتی، آخرین مدرک تحصیلی همسر، تناسب میانگین درآمد با فعالیت، بیماری خاص جسمی یا روان شناختی، وضعیت شیفت کاری در اورژانس، شیفت روز تعطیل در یک سال گذشته و تعداد شیفت شب در یک سال گذشته اختلاف معناداری ندارد (جدول ۲).

جدول ۲) گزارش رابطه بین نمرهٔ پرسشنامهٔ امید به زندگی و ویژگیهای جمعیتشناختی و شغلی افراد

P value	انحراف معیار	میانگین نمره پرسشنامهٔ امید به زندگی	وضعيت	متغير	رديف
۰/۲۲		۱۶۰/۸۵	زن		
o/ Y F	W5/99	184/14	مرد	جنسیت	١
, ,	۱۸/۸۲	140/90	مجرد	, f.,	.,
<0/0001	WW/8Y	151/44	متاهل	تأهل	۲
	۳۸/۲۰	17°/47	بلی	I	٣
<0/0001	19/09	۱۵۲/۳۲	خير	فرزند داری	۲
. / \	۳۸/۲۰	14°/44	بلی	He	k
<0/0001	18/74	144/89	خير	زندگی کردن فرزند با والد مورد سؤال	۲
	۲۹/۹۵	۱۳۵/۷۶	پزشک عمومی		
<0/0001	۲۳/۲۹	174/4°	متخصص طب اورژانس	رشته تحصيلي	۵
	۳۴/۸۷	۱۵۵/۶۸	متخصص در رشته های دیگر		
, ,	WF/17	۱۶۸/۸۵	بلی	He . Late	
∘/∘∘∧	٧/٣۶	۱۴۸/۳۵	خير	شاغلبودن همسر فرد مورد سؤال	۶
	۳۷/۵۶	184/90	دكترا		
۰/۵۶	9/٧۴	۱۷۲/۰۸	فوق ليسانس	آخرین مدرک تحصیلی همسر	٧
	۲۰/۷۴	184/71	ليسانس		
	Y9/YY	109/FY	بلی	W . 1 7 6.1 1	
۰/۶۹	۳۱/۹۰	181/48	خير	تناسب میانگین درآمد با فعالیت	٨
	۳۶/۷۱	145/14	خیلی کم		
	۱۷/۳۰	1ΥΥ/ΔΥ	کم		
<0/0001	۳۱/۲۸	184/99	 متوسط	میزان رضایت از وضعیت کاری خود	٩
	•	•	زیاد		
	•	•	 بسیار زیاد		
	۳۸/۹۰	۱۳۵/۱۴	خیلی کم		
	17/40	۱۵۵/۹۸	کم		
<0/0001	۲۱/۸۸	188/YI	 متوسط	میزان رضایت از وضعیت زندگی خود	1.
	1Δ/•Α	Y1\%/YY	زیاد		
	•	•	بسیار زیاد		
	Y ₀ /YY	1ΔΥ/1Υ	خیلی کم		
	*Y/Y°	۱۸۸/۵۰	کم		
<0/0001	۲۰/۷۵	181/90		تمایل به تغییر حرفه در وضعیت فعلی	11
	Y1/18	۱۷۱/۵۵	 زیاد		
	۳۶/۸۱	140/14	بسیار زیاد		
	۳۸/۹۸	140/01			
		184/48	کم کم		
<0/0001	۲۱/۹۵	154/04	 متوسط	مسافرت جهت رهایی از استرس شغلی	١٢
		۱۲۹/۵۸	زیاد		
	0/00	191/00	 بسیار زیاد		

وره ۱۳، شماره ۱، ۱۴۰۳ نشریه علمی پژوهشی طب انتظامی

بررسی توصیفی تحلیلی عوامل مؤثر بر امید به زندگی در پزشکان نوبتکار بخش اورژانس

P value	انحراف معیار	میانگین نمره پرسشنامهٔ امید به زندگی	وضعيت	متغير	ردیف
	۵۰/۰۲	140/94	خیلی کم		
	T8/Y°	۱۷۲/۵۶	کم		
<0/0001	YW/9Y	105/04	متوسط	مراوده با نزدیکان جهت رهایی از استرس شغلی	١٣
	YY/2Y	184/18	زیاد		
	•	0	بسیار زیاد		
	kk/1 k	140/44	خیلی کم		
	۲۱/۳۸	180/88	 کم		
<0/0001	~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~~	۱۷۵/۹۸	متوسط	ورزش جهت رهایی از استرس شغلی	16
	۱۸/۲۵	179/191	زیاد		
	•/••	141/00	بسیار زیاد		
	k4\kk	۱۱۲/۱۵	به هیچ وجه		
	۲۵/۰۶	۱۵۲/۷۸	تاحدودی		
<0/0001	Y8/Y8	175/18	نسبتاً	خواب مناسب در طی شبانه روز	۱۵
	YY/19	188/18	اغلب		
	•	۱۷۵/۰۰	کاملا		
	YY/9.A	184/44	بلی	all a land	
۰/۱۶	mm/9k	109/80	خیر خیر	بیماری خاص جسمی یا روانشناختی	18
	#k\#k	188/29	بلی		
۰/۰۴	Y9/88	۱۵۸/۷۴	خیر	بیماری خاص اعضای درجهٔ اول خانواده	17
	۳۰/۹۶	184/40	شيفت ثابت		
۰/۲۲	۳۱/۲۸	181/18	 شیفت در گردش	وضعیت نوبت کاری در اورژانس	١٨
	18/40	۱۷۲/۸۴	صبح		
	•	۱۷۵/۰۰	عصر		
<0/0001		14°/01	 شب	زمان نوبت کاری در اورژانس	19
	۲۸/۱۹	184/14	 ترکیبی شیفت صبح، عصر و شب		
	۳۱/۹۵	181/80	بلی		.,
۰/۵۵	۸/۱۳	ነ۵۴/۸۸	خیر	شیفت روز تعطیل در یک سال گذشته	۲۰
	۲۵/۵۷	188/49	کمتر از ۴ شب		
۰/۱۸۳	YY/11	184/18	۵ تا ۸ شب	تعداد شیفت شب در یک سال گذشته	۲۱
	۴۱/۵۰	۱۵۶/۸۵	 بیشتر از ۸ شب		

از رگرسیون خطی چندگانیه و گام بیه گام، بیرای آزماییش اینکه آیا متغیرهای سن، مدت زمان تأهل، تعداد فرزنید، سابقه کار کلی و سابقهٔ کار در اورژانیس بیه طور قابل توجهی نمیرهٔ پرسشیامهٔ امیید بیه زندگی را پیشبینی میکننید، استفاده شد. نتاییج تست مدل رگرسیون بیرازش شده (میدل اوّل) عبارت بود از:

نمرهٔ پرسشنامهٔ امیید به زندگی = ۲۹۴/۹۱ - ۴/۸۴ (سین) + $^{\circ}$ /۱/۵۰ (سین) + $^{\circ}$ /۱/۵۰ (سین) + $^{\circ}$ /۱/۵۰ (سیابقه کار کلی) + $^{\circ}$ /۱/۵۰ (سیابقه کار در اورژانس) رگرسیون کلی از نظر آماری معنی دار بود (۲۱۶–82 و $^{\circ}$ /۲/۵۰ و $^{\circ}$ /۱/۵۰ و $^{\circ}$ /۱/۵۰ و $^{\circ}$ /۱/۵۰ ($^{\circ}$ /۱/۵۰)

رگرسیون گام به گام، با استفاده از حذف به عقب (مدف به عقب p<-۱/۱۰) با آستانه حذف p<-۱/۱۰ انجام شد. طبق نتایج رگرسیون، سن پزشکان، مدت زمان تأهل، سابقهٔ کار کلی و سابقهٔ کار در اورژانس، به طور قابل توجهی نمرهٔ پرسشنامهٔ امید به زندگی را پیشبینی میکنند. بنابرایی متغیرهای اشاره شده وارد مدل دوم رگرسیونی شده و تعداد فرزندان از این مدل خارج شد. (جدول ۳)

نتایی رگرسیون مشخص کرد که سن پزشکان، به طور قابل توجهی نمرهٔ پرسشنامهٔ امیید به زندگی را پیشربینی میکنید ($(-2)^4$ - $(-2)^4$). درحالی که، میدت زمان تأهل پزشکان، به طور قابل توجهی نمرهٔ پرسشنامهٔ امیید به زندگی را پیشربینی میکنید ($(-2)^4$) و $(-2)^4$ و $(-2)^4$


در ادامه، نمودارهای مربوط به پراکندگی (شکل ۲) و نمودار P-P، برای باقیماندهٔ استانداردشدهٔ رگرسیون (شکل ۳) و نمودار هیستوگرام (شکل ۴) آورده شده است. همچنین همبستگی پیرسون، برای بررسی قدرت رابطه خطی دو به دو، بین متغیرهای فوق استفاده شد که نتایج آن در جدول ۴ آورده شده است.

طلوعی و همکاران


р	Т	Beta	SE	В	متغیرهای پیشبین واردشده	مدل
0/0001	14/128	-	Y1/44A	249/910	مقدار ثابت	
0/0001	-5/494	-1/°YY	°/YFY	-4/7/61	سن	
0/001	٣/٢۶١	۰/۳۶۲	0/481	1/0.4	مدت زمان تأهل	
۰/۲۱۷	1/۲۳۸	0/109	٣/٣٢۵	4/110	تعداد فرزند	1
0/0001	4/1/4	0/844	۰/۶۷۳	۲/ ۸۳۶	سابقهٔ کار کلی	
۰/۰۱	4/094	۰/۲۱۵	0/481	1/190	سابقهٔ کار در اورژانس	
		F	R2=، ۱۹۸ ،R2	توجه: R=۰/۴۶۵، ۲۱۶/۰		
0/0001	۱۴/۲۴۵	-	۲۱/۰۵۸	Y99/9AY	مقدار ثابت	
0/0001	-819014	-1/171	۰٬۷۳۱	-0/040	سن	
0/001	4/084	°\&h.h.	°/46k	1/494	مدت زمان تأهل	
0/0001	۴/۶۲۰	۰/۶۶۵	∘/۶۵۶	٣/٠٢٩	تعداد فرزند	۲
۰/۰۰۶	Y/ YY 1	۰/۲۲۸	۰/۴۵ ۷	1/484	سابقهٔ کار کلی	
0/0001	14/440	-	Y1/00A	Y99/9AY	سابقهٔ کار در اورژانس	
		1	R2=، ۱۹۶، R2=،	توجه: R=0/۴۵۹، ۲۱۱،		


جدول ۴) نتایج همبستگی پیرسون (Pearson correlation coefficient)

سابقهٔ کار در اورژانس	سابقهٔ کار کلی	تعداد فرزند	مدت زمان تأهل	سن	نمرهٔ پرسشنامهٔ امید به زندگی	متغيرها
۰/۰۸۴	۰/۱۰۴	۰/۲۳۶	o/1YY	-0/041	1/000	نمرهٔ پرسشنامهٔ امید به زندگی
۰/۶۸۸	۰/٩٠٣	۰/۵۲۸	۰/۲۵۸	1/000	-o/o ۴ 1	سن
۰/۴۸۱	۰/۲۲۹	۰/۶۹۵	1/000	۰/۲۵۸	o/\YY	مدت زمان تأهل
۰/۴۰۰	۰/۵۲۸	1/000	۰/۶۹۵	۰/۵۲۸	۰/۲۳۶	تعداد فرزند
۰/۵۹۶	1/000	۰/۵۲۸	۰/۲۲۹	۰/٩٠٣	۰/۱۰۴	سابقهٔ کار کلی
1/000	∘/۵۹۶	۰/۴۰۰	۰/۴۸۱	۰/۶۶۸	۰/۰۸۴	سابقهٔ کار در اورژانس



شکل ۲) نمودار پراکندگی برای باقیماندهٔ استانداردشدهٔ رگرسیون برای پیشبینی نمرهٔ پرسشنامهٔ امید به زندگی






شکل p) شکل p-p نرمال برای باقیماندهٔ استانداردشدهٔ رگرسیون برای پیشبینی نمرهٔ پرسشنامهٔ امید به زندگی

دوره ۱۳، شماره ۱، ۱۳۰۳ - نـشـریــه علمی پژوهشی طـب انـتـظـامــی

بررسی توصیفی تحلیلی عوامل مؤثر بر امید به زندگی در پزشکان نوبتکار بخش اورژانس



شکل ۴) هیستوگرام باقیماندهٔ استانداردشدهٔ رگرسیون برای پیشبینی نمرهٔ پرسشنامهٔ امید به زندگی

بحث

ایـن پژوهـش بـا هـدف بررسـی عوامـل مؤثـر بـر امیـد بـه زندگی در پزشـکان شـیفت کار شـاغل در بخـش اورژانـس انجام شـد.

تغییرات دهههای اخیر، سیستم کار و اقتصاد را دستخوش تغییراتی کرده است. هر چند شیفت کاری در بیـن کادر درمانـی از ابتـدا مرسـوم بـوده، ولـی کمبـود نیـروی کار با تجربه از یکسو و نیازهای اقتصادی افراد از سوی دیگر، این موضوع را بیش از پیش در این قشر مورد توجه قرار داده است. در این تحقیق، برخلاف تحقیقات پیشین که عمدتاً معطوف به پرستاران بود، عوامل مؤثر بر امید به زندگی و مشکلات شیفت کاری در پزشکان عمومی و متخصص شاغل در بخش اورژانس مراکز درمانی دولتی و خصوصی شهر تهران بررسی شد. مراکز درمانی به عنوان سازمان فراهم کننده خدمات درمانی برای شهروندان، باید تمهیداتی در جهت ارتقاء سطح کیفیت و امید به زندگی و حـل مشـكلات ناشـي از موقعيـت شـغلي كاركنـان بالينـي (کادر درمان) در نظر بگیرد، زیرا با داشتن کارکنان پویا، با انگیزه، امیدوار و هدفمند، بهتر و سریعتر میتواند به اهداف بلندمدت سازمانی خود برسد و خدمات بهینه به مراجعه کننـدگان ارائـه دهـد.

پژوهشهای قبلی، تأثیرات قابل توجه نوبت کاری را بر خواب، خلق افسرده و اضطراب، مصرف مواد، اختلال در شناخت، کیفیت پایین زندگی و حتی افکار خودکشی روشن ساختهاند که اختلالات خواب واضح اغلب زمینه ساز عواقب سلامت روان شیفت کاری ذکرشدهاست [۲۰].

اصغـری و همـکاران (۱۳۹۲)، در مطالعـهای بـر روی ۱۵۰ نفـر از کارگـران شـاغل در یکـی از صنایـع خودروسـازی شـهر تهـران، نشـان دادهانـد کـه بالاتریـن میـزان شـیوع مشـکلات ناشـی از نوبـت کاری مربـوط بـه اختـلال در زندگـی خانوادگـی، فـردی و اجتماعـی نوبـت کاران اسـت [۲۱].

یافته های مطالعه مهدیزاده نشان داده است که پرستارانی که تعداد شیفت شب بیشتری در برنامه کاری خود داشتهاند و کسانی که ساعات خواب کمتر از ۲ ساعت

در طبی شب کاری داشته اند، همچنین پرستارانی که در تنظیم برنامهٔ کاری خود مشارکت نداشته اند و پرسنل شاغل در بخشهای ICU، با تغییرات هورمونی بیشتری مواجه هستند [۲۲]. در مطالعه کاکویی، مقایسه مدل تغییرات زمانی ترشح ملاتونین در پرستاران نوبت کار و ثابت کار، حاکی از این واقعیت است که نوبت کاری به طور سرشتی برای انسان غیرطبیعی و غیرعادی بوده، با تغییر الگوی ترشح ملاتونین (کاهش قابل توجه میزان ترشح ملاتونین در نقطه اوج یعنی ساعت ۴ صبح و اوایل صبح یعنی ساعت ۷ به طور قطع چرخه خواب، میزان هوشیاری در شب و فعالیت روزانه برهم خواهد خورد [۳۲]. شواهدی وجود دارد مبنی بر اینکه پرستاری حرفهای پر استرس است ۲۲].

در این پژوهش، ارتباط سن، وضعیت تأهل، صاحب فرزند، زندگی کردن فرزند با والد مورد سؤال، رشتهٔ تحصیلی، شاغلبودن همسر فرد مورد سؤال، آخرین مدرک تحصیلی، تناسب میانگین درآمد با فعالیت، میزان رضایت از وضعیت کاری خود، میزان رضایت از وضعیت زندگی خود، تمایل به تغییر حرفه در وضعیت فعلی، مسافرت برای رهایی از استرس کاری، مراوده با نزدیکان برای رهایی از استرس کاری، ورزش برای رهایی از استرس کاری، خـواب مناسـب در طـی شـبانهروز، بیمـاری خاصـی جسـمی یا روان شناختی، بیماری خاص اعضای درجهٔ اول خانواده، شیفت روز تعطیل در یک سال گذشته و تعداد شیفت شب دریک سال گذشته با وضعیت امید به زندگی مورد بررسی قـرار گرفـت کـه براسـاس نتایـج بـه دسـت آمـده، نمـرهٔ آزمـون امید به زندگی با وضعیت تأهل، فرزندداری، زندگی فرزند با والد مورد سؤال، رشتهٔ تحصیلی، شاغلبودن همسر، زندگی کردن فرزند با والد مورد سؤال، رشتهٔ تحصیلی، شاغلبودن همسـر فـرد مـورد سـؤال، ميـزان رضايـت از وضعیت کاری خود، میزان رضایت از وضعیت زندگی خود، تمایل به تغییر حرفه در وضعیت فعلی، مسافرت برای رهایی از استرس شغلی، مراوده با نزدیکان برای رهایی از

استرس شغلی، ورزش برای رهایی از استرس شغلی، خواب مناسب در طی شبانه روز، بیماری خاص اعضای درجه اول خانـواده و وضعیـت شـیفت در اورژانـس ارتبـاط معنـاداری داشت. همچنین، سن پزشکان، مدت زمان تأهل پزشکان، سابقهٔ کار کلی پزشکان و سابقهٔ کار در اورژانس پزشکان، به طور قابل توجهی نمرهٔ آزمون امید به زندگی را پیشبینی کرد. نوع شیفت کاری با امید به زندگی رابطه معناداری داشت، به این معنی که در افرادی که عمدتاً شیفت صبح یا عصر را بر عهده داشتند، نمرهٔ میزان امید به زندگی آنها نسبت به افراد در شیفتهای در گردش و شب بالاتر بود. این نتایج، با پژوهشهای میرمحمدی و همکاران (۱۳۹۰)، *اردکانی* و همکاران (۱۳۸۶)، Ferri و همکاران (۲۰۱۶) و Barrau-Baumstarck و همـكاران (۲۰۰۹) همسـو اسـت [۲۷-۲۴]. محققین فوق معتقدنید شیفت کاری نامناسب به دلیل اثر منفی بر سلامت اجتماعی، سبب می شود تا فرد به واسطه نداشتن برنامهٔ زندگی مدون و از پیش مشخص، از تعاملات اجتماعی منظم و پیوسته محروم گردد.

یافته های پژوهش ما، اختلاف معناداری در نمرهٔ آزمون امید به زندگی پزشکان در گروههای مختلف براساس شیفت روز تعطیل در یک سال گذشته و تعداد شیفتهای شب پیدا نکرد. این نتیجه با یافته های پژوهشهای قبلی، از جمله کارهای فرهناکی و همکاران (۱۳۹۳)، سبکرو فرهماکاران (۱۳۹۳)، سبکرو شبانه و تعطیلات رسمی بر تأثیر منفی شیفت کاری شبانه و تعطیلات رسمی بر زندگی خانوادگی غیرهم سو است. همچنین اگرچه نتایج فرهناکی و همکاران (۱۳۹۳) و مرز تعطیل در منزل و کیفیت نامناسب زمان حضور، به و روز تعطیل در منزل و کیفیت نامناسب زمان حضور، به جهت عدم توانایی نظارت دقیق بر امورات منزل به واسطه خستگی ناشی از شیفت کاری اشاره دارد، ولی مطالعهٔ منفی با امید به زندگی در شیقت کاران شب و روزهای تعطیل نیافت [۲۱، ۲۸، ۲۸].

Barrau-Baumstarck (۲۰۱۰)، Barnes-Farrell و همـکاران (۲۰۰۹) معتقدنـد افـراد شـیفت کار بایـد سـعی در کنارآمـدن و سـازش بـا شـرایط شـغلی خـود داشـته باشـند؛ زیـرا در صـورت عـدم سـازش بـا شـرایط توانایـی مقابلـه بـا مشـکلات پیـش آمـده در زمینههـای فـردی، خانوادگـی و شـغلی را نخواهنـد داشـت و ایـن امـر موجبات فرسـودگی شخص شـیفت کار را فراهـم مــیآورد [۳۰٬۲۷].

مطالعـهٔ مـا محدودیتهایـی داشـت کـه از جملـه میتـوان بـه وسـعت و چندعاملیبـودن مـوارد تأثیرگـذار بـر امیـد بـه زندگـی اشـاره نمـود کـه احتمـال تحـت تأثیـر قـرار گرفتـن نتایــج مطالعـه بـا ایــن موضوعـات وجـود دارد. لازم است تـا در مطالعـات تکمیلـی آینـده، تأثیـر متغیرهایـی ماننـد وضعیـت اسـتخدامی و رژیـم غذایـی نیـز بـر امیـد بـه زندگـی مـورد بررسـی قـرار گیـرد. محدودیـت دیگـر ایــن پژوهـش در محـود بهطوری کـه پاسـخدهندگان بـه جمـعآوری دادههـا بـود، بهطوری کـه پاسـخدهندگان بـه واسـطهٔ خودسـنجی ممکـن اسـت در پاسـخدادن سـوگیری

داشته باشند.

نتیجه گیری

نتایج حاصل از این مطالعه مشخص کرد که شیفت کاری نامناسب میتواند موجب اختلال در زندگی فردی، خانوادگی و شغلی پرسنل کادر درمان شود. همچنین عوامل مختلفی از جمله ویژگیهای جمعیت شناختی، شیفت کاری، درآمد، خواب، وضعیت زندگی، مسافرت، ورزش، مراوده با نزدیکان، بیماری خاص جسمی یا روانشناختی و بیماری خاص اعضای خانواده، بر امید به زندگی پزشکان شیفتکار اشت.

به نظرمیرسد کاهش ساعات کاری پرسنل، اجتناب از شیفتهای ثابت شب و یا شیفتهای بیش از حد روزهای تعطیل رسمی در بین پزشکان شاغل در بخش اورژانس، میتواند از فرسودگی شغلی و کاهش امید به زندگی جلوگیری نماید. در این صورت، سرمایهٔ اجتماعی پرسنل بالات رفته و رضایت شغلی و کیفیت مراقبت از بیماران نیز افزایش میابد. با در نظر داشتن ذات حساس شغل پزشکی و به ویژه، پزشکان شاغل در بخش اورژانس و نیاز به حضور شبانهروزی در بخش درمان کشور، لازم است تا تدابیری جهت حفظ سلامتی این گروه از پزشکان از جمله مشوقهای جایگزین اندیشیده شود.

نكات بالينى كاربردى براى پليس:

بر اساس یافتههای این مطالعه و با توجه به اهمیت موضوع امید به زندگی و شیفتهای کاری پزشکان، نکات بالینی و پزشکی زیر میتواند برای جوامع انتظامی و پلیس نیز کاربرد داشته باشد که در زیر به برخی از آنها اشاره می گردد:

۱. توجه به سه سلامت روحی و روانی پزشکان. به دلیل استرس بالا و شرایط دشوار شغلی، ضروری است که جوامع انتظامی از برنامههایی برای حمایت روانی پزشکان اورژانس استفاده کنند. فراهم آوردن خدمات مشاورهای و روانشناسی میتواند به کاهش استرس و افزایش امید به زندگی آنها کمک کند.

۲. افزایـش آگاهـی در مـورد اهمیـت شـیفتهای کاری. بـا توجـه بـه ارتبـاط معنـادار نـوع شـیفت کاری بـا امیـد بـه زندگـی پزشـکان، جوامـع انتظامـی بایـد آگاهـی بیشـتری در مـورد اثـرات مضـر شـیفتهای طولانـی و شـبانه بـر سـلامت جسـمی و روانـی داشـته و در سیاسـتگذاریها و اسـتخدامها بـه آن توجـه کننـد.

۳. ترویج فعالیتهای ورزشی و تفریحی. تشویق پزشکان به مشارکت در فعالیتهای ورزشی و تفریحی میتواند به کاهش استرس و بهبود کیفیت خواب و زندگی آنها کمک کند. جوامع انتظامی میتوانند با حمایت از برنامههای ورزشی و تفریحی برای کارکنان به بهبود وضعیت سلامت آنها یاری برسانند.

بررسی توصیفی تحلیلی عوامل مؤثر بر امید به زندگی در یزشکان نوبت کار بخش اورژانس

زندگی پزشکان را بهبود بخشند و در نتیجه به بهبود خدمات درمانی ارائه شده به جامعه نیز کمک نمایند.

سهم نویسندگان: حامد اقدم، امید شیرزاد و پویا وکیلی پور؛ ارائه ایده و طراحی مطالعه، امید شیرزاد؛ جمع آوری دادهها، شبنم طلوعی؛ تجزیه و تحلیل دادهها. همه نویسندگان در نگارش اولیه مقاله و بازنگری آن سهیم بودند و همه با تأیید نهایی مقاله حاضر، مسئولیت دقت و صحت مطالب مندرج در آن را می نیرند.

تعارض منافع: بدینوسیله نویسندگان مقاله تصریح مینمایند که هیچگونه تعارض منافعی در قبال مطالعه حاضر وجود ندارد.

حرب . حامــی مالــی: در ایــن مقالــه هیچگونــه حامــی مالــی وجــود داشت. ۴. ایجاد حمایتهای خانوادگی. با توجه به تأثیر وضعیت خانواده بر امید به زندگی پزشکان، جوامع انتظامی میتوانند با ایجاد برنامههایی برای بهبود زندگی خانوادگی پزشکان، نظیر خدمات مشاورهای خانوادگی، نقش موثری در افزایش کیفیت زندگی آنان ایفا کنند.

۵. تعییان شایفتهای حمایتی. با توجه به یافتهای جوامع انتظامی باید اقداماتی در راستای تغییر شایفتهای کاری پزشکان، از جمله کاهش شایفتهای شابانه و طولانی را مدنظر قارا دهند. ایان تغییارات میتواند به کاهش فرسودگی شغلی و بهبود کیفیات مراقبت از بیماران کمک کند.

این نکات میتواند به جوامع انتظامی کمک کند تا با برنامهریزی بهتر و حمایتهای لازم، کیفیت

Reference

- Kakooei H, Zamanian Ardakani Z, Karimian M, Ayttollahi T. Twenty four hours circadian cortisol profile in shift work nurses. Armaghane danesh. 2009;14(1):47-56. http://armaghanj.yums.ac.ir/ article-1-571-en.html
- Fesharaki GM, Rozati M, Tanhai A. The longitudinal study of the relationship between work shift and blood pressure in workers of Mobarakeh Steel Company of Isfahan in 2007-2009. Arak Med Univ J. 2010;13(4):68-74. http://jams.arakmu.ac.ir/article-1-587-en.html
- Choobineh A, Soltanzadeh A, Tabatabaee S, Jahangiri M, Khavaji S. Shift work-related health problems in" 12-hour shift" schedule in petrochemical industries. Iran Occup Health. 2010;7(1):44-53. https://ijoh.tums.ac.ir/index.php/ijoh/article/view/32
- Fischer FM. What do petrochemical workers, healthcare workers, and truck drivers have in common? Evaluation of sleep and alertness in Brazilian shiftworkers. Cad Saúde Pública. 2004;20:1732-8. https://doi.org/10.1590/s0102-311x2004000600033
- Ha M, Park J. Shiftwork and metabolic risk factors of cardiovascular disease. J Occup Health. 2005;47(2):89-95. https://doi.org/10.1539/joh.47.89
- Costa G. Shift work and occupational medicine: an overview. Occup Med. 2003 53(2):83-8. https:// doi.org/10.1093/occmed/kqg045
- 7. Pires MLN, Teixeira CW, Esteves AM, Bittencourt LRA, Silva RS, Santos RF, et al. Sleep, ageing and night work. Brazilian J Med Biol. 2009;42:839-43. https://doi.org/10.1590/S0100-879X2009005000011
- 8. Kilpatrick K, Lavoie-Tremblay M. Shiftwork: what health care managers need to know. Health Care Manag. 2006;25(2):160-6. https://doi.org/10.1097/00126450-200604000-00009
- Dehghan, A. A. S., Nejad, Z. C., & Maznab, A. A. (2019). The Effectiveness of Positive teaching with an Islamic Approach in a group way in Nurses' Life

- Expectancy. Journal of Nursing Education (JNE), 8(4).
- Snyder CR. The psychology of hope: You can get there from here: Simon and Schuster; 1994. https://books.google.com/books/about/The_Psychology_of_Hope.html?id=b3bz8tzBvC0C
- Carr A. Positive psychology: The science of happiness and human strengths. Routledge; 2013. https://doi.org/10.4324/9780203506035
- 12. Snyder CR, Harris C, Anderson JR, Holleran SA, Irving LM, Sigmon ST, et al. The will and the ways: development and validation of an individual-differences measure of hope. J Personal Soc Psychol. 1991;60(4):570. https://psycnet.apa.org/doi/10.1037/0022-3514.60.4.570
- 13. Aburto JM, Villavicencio F, Basellini U, Kjærgaard S, Vaupel JW. Dynamics of life expectancy and life span equality. PNAS. 2020;117(10):5250-9. https://doi.org/10.1073/pnas.1915884117
- 14. Goto A, Takao T, Yoshida Y, Kawazu S, Iwamoto Y, Terauchi Y. Causes of death and estimated life expectancy among people with diabetes: A retrospective cohort study in a diabetes clinic. J Diabetes Investig. 2020;11(1):52-4. https://doi.org/10.1111/jdi.13077
- 15. Cao W, Qi X, Yao T, Han X, Feng X. How doctors communicate the initial diagnosis of cancer matters: cancer disclosure and its relationship with Patients' hope and trust. Psychooncology. 2017;26(5):640-8. https://doi.org/10.1002/pon.4063
- Clayton JM, Hancock K, Parker S, Butow PN, Walder S, Carrick S, et al. Sustaining hope when communicating with terminally ill patients and their families: a systematic review. Psychooncology. 2008;17(7):641-59. https://doi.org/10.1002/pon.1288
- Miller JF, Powers MJ. Development of an instrument to measure hope. Nurs Res. 1988;37(1):6-10. https://pubmed.ncbi.nlm.nih.gov/3340583/
- 18. Darvishi H. Study the function of mental imagina-

tion and relaxation on reduction of stress, depression and increasing life expectancy in female patients with breast cancer,(2009), Ahwaz golestan Hospital: MA thesis, IAU, Ahwaz Branch. https://www.sid.ir/paper/189654/en

- Brown JP, Martin D, Nagaria Z, Verceles AC, Jobe SL, Wickwire EM. Mental health consequences of shift work: an updated review. Curr Psychiatry Rep. 2020;22:1-7. https://doi.org/10.1007/s11920-020-1131-z
- Asghari M, Ahmadnezhad I, Rahmani A, Sadeghi A, Abbassinia M, Rezaie E, Ahsani H. Investigation of disorders and problems caused by shift work in an automotive industry. J Univ Med Sci. 2013. http:// journal.nkums.ac.ir/article-1-25-en.html
- 21. Mahdizadeh SM. Assessing the level of serum Cortisol and Aldestrone of nursing personal in various work shifts in Mashhad medical sciences hospitals. Iran Occupational Health. 2009;6(1):56-60. http://ioh.iums.ac.ir/article-1-170-en.html
- Kakooei H, Zamanian Ardakani Z, Karimian SM, Ayattollahi SN. Twenty Four-hour circadian melatonin profile among women shift work nurses. J Adv Med Biomed Res. 2009;17(68):75-84. http:// journal.zums.ac.ir/article-1-993-en.html
- Mirmohammadi S, Mehrparvar A, Kamali Z, Mostaghaci M. Evaluation or the relationship between shift work and sleepiness in nurses. 2011. http:// tkj.ssu.ac.ir/article-1-43-en.html
- 24. Ardekani ZZ, Kakooei H, Ayattollahi S, Choobineh A, Seraji GN. Prevalence of mental disorders among shift work hospital nurses in Shiraz, Iran. Pak J Biol Sci. 2008;11(12):1605-9. https://doi.org/10.3923/pjbs.2008.1605.1609
- 25. Ferri P, Guadi M, Marcheselli L, Balduzzi S, Magnani D, Di Lorenzo R. The impact of shift work on the psychological and physical health of nurses in a general hospital: a comparison between rotating night shifts and day shifts. Risk Manag Healthc Policy. 2016:203-11. https://doi.org/10.2147/rmhp.s115326
- Barrau-Baumstarck K, Rebeschini E, Dalivoust G, Durand-Bruguerolle D, Gazazian G, Martin F. Shiftwork and quality of life among critical care nurses and paramedical personnel. Presse Med. 2009;38(3):346-53. https://doi.org/10.1016/j.lpm.2008.06.018
- 27. Farahnaki Z, Pournajaf A, Karami A, Abbasi A, Karchani M. Investigation of shift work disorders in nurses working at teaching hospitals of Ilam. Hospital. 2014;13(1):35-42. http://jhosp.tums.ac.ir/article-1-5290-en.html
- Sabokroo M, Kalhorian R, Kamjoo Z, Taleghani G. Work-family Conflict: The role of organizational supportive perception in turnover intention (Case study of nurses of Tehran's hospitals. J Public Administration. 2011;3(6):111-236. https://dor.isc. ac/dor/20.1001.1.20085877.1390.3.6.7.6
- 29. Barnes-Farrell JL, Davies-Schrils K, McGonagle A, Walsh B, Di Milia L, Fischer FM, et al. What as-

pects of shiftwork influence off-shift well-being of healthcare workers? Appl Ergon. 2008;39(5):589-96. https://doi.org/10.1016/j.apergo.2008.02.019